

Some properties of q -perfect matroid designs

KAWABUCHI, Shinya

Kumamoto University

Joint work with SHIROMOTO, Keisuke

Matroid is a generalization of linear independence.

• Matroid

- Combinatorial structure based on the concept of the linear independence.

• q -Matroid

- q -analogue of a matroid
- R. Jurrius, R. Pellikaan established. (2018)

q -analogue: the way of generalization

	normal	q -analogue
finite set	$[0, 1, \dots, n]$	\mathbb{F}_q^n
size of X	$ X $	$\dim X$
subset	subset	subspace
union	$X \cup Y$	$X + Y$
intersection	$X \cap Y$	$X \cap Y$

Preliminaries

- q : a prime power
- $E(= \mathbb{F}_q^n)$: an n -dimensional vector space over \mathbb{F}_q
- $\mathcal{L}(X)$: the collection of all subspaces of a vector space X
- $\begin{bmatrix} X \\ k \end{bmatrix}_q$: the collection of all k -dimensional subspaces of a vector space X

Definition [q -matroid]

A q -matroid is a pair (E, r) satisfying (qR1), (qR2) and (qR3):

	q -matroid	matroid
ground space E	\mathbb{F}_q^n	$[n] = \{1, 2, \dots, n\}$
rank function r	$\mathcal{L}(E) \rightarrow \mathbb{Z}_{\geq 0}$	$2^E \rightarrow \mathbb{Z}_{\geq 0}$
1 st axiom	(qR1) $0 \leq r(A) \leq \dim A$	(R1) $0 \leq r(A) \leq A $
2 nd axiom	(qR2) $A \subseteq B \Rightarrow r(A) \leq r(B)$	(R2) $A \subseteq B \Rightarrow r(A) \leq r(B)$
3 rd axiom	(qR3) $r(A + B) + r(A \cap B) \leq r(A) + r(B)$	(R3) $r(A \cup B) + r(A \cap B) \leq r(A) + r(B)$

- integer $k : 0 \leq k \leq n$

Example [The uniform q-matroid]

$$r(X) := \min\{\dim X, k\}.$$

Then, the pair (E, r) is a q -matroid.

Remark. This q -matroid is called a ***uniform q-matroid*** $U_{k,n}[\mathbb{F}_q]$

- $M = (E, r)$: q -matroid

Definition [Flat of a q -matroid]

$F \subseteq E$ is a *flat* of M if and only if

$$x \in \begin{bmatrix} E \\ 1 \end{bmatrix}_q \setminus \begin{bmatrix} F \\ 1 \end{bmatrix}_q \Rightarrow r(F + x) = r(F) + 1$$

holds.

- $M = (E, r)$: matroid

Definition [Flat of a matroid]

$F \subseteq E$ is a *flat* of M if and only if


$$x \in E - F \Rightarrow r(F \cup x) = r(F) + 1$$

holds.

Remark

We denote \mathcal{F} as a collection of all flats in M .

- If a flat F satisfies $r(F) = i$, F is called **i -flat**.
- The collection of all i -flats is denoted by \mathcal{F}_i .




Proposition [axiom of flats of q -matroid] (E. Byrn et al. 2022 +)

\mathcal{F} is a collection of all flats of q -matroid
 $\Rightarrow \mathcal{F}$ satisfies (qF1), (qF2) and (qF3)

- $F_1, F_2, F \in \mathcal{F}$

(E, r)	q -matroid	matroid
1 st axiom	(qF1) $E \in \mathcal{F}$	(F1) $E \in \mathcal{F}$
2 nd axiom	(qF2) $F_1 \cap F_2 \in \mathcal{F}$	(F2) $F_1 \cap F_2 \in \mathcal{F}$
3 rd axiom	(qF3) $\exists! F' \in \mathcal{F}_{r(F)+1} \text{ s.t. } F + x \leq F'$ $\left(\forall x \in \begin{bmatrix} E \\ 1 \end{bmatrix}_q \setminus \begin{bmatrix} F \\ 1 \end{bmatrix}_q \right)$	(F3) $\exists! F' \in \mathcal{F}_{r(F)+1} \text{ s.t. } F \cup x \subseteq F'$ $(\forall x \in E - F)$

Hasse diagram of flats



t -dimensional subspace included in λ blocks

- $\mathcal{B} \subseteq \binom{E}{k}_q$

Definition [subspace design]

A t -($n, k, \lambda; q$) subspace design is a pair (E, \mathcal{B}) with the property that every t -dimensional subspace of E is contained in exactly λ elements of \mathcal{B} .

Remark

- A member of \mathcal{B} is called a ***block***.
- A subspace design t -($n, k, 1; q$) is called a ***q-Steiner system*** denoted by $\mathcal{S}(t, k, n; q)$.
- Subspace designs have been actively studied because of their application to random network coding.

Perfect matroid designs have a lot of t -designs.

- **Perfect matroid design (PMD)**

- A matroid whose flats of the same rank all are the same cardinality.
- U.S.R. Murty, P. Young and J. Edmonds established(1970).
- M. Deza and N.M Singh studied some properties of PMD and the PMD of rank 4 .
- There are many kinds of blocks of t -design in PMD.
 - flats, bases, circuit
- **Steiner systems induce PMDs**

- **q -perfect matroid design (q -PMD)**

- q -analogue of PMDs
- E. Byrne, M. Ceria, S. Ionica and R. Jurius (2022)
- **q -Seiner systems induce q -PMDs**

	PMD	q -PMD
How to construct non trivial (q)-PMD	<ul style="list-style-type: none"> • Projective geometries • Affine geometries • Affine triple systems • Steiner system 	<ul style="list-style-type: none"> • Steiner system • ??? • ??? • ???
α -sequence	the cardinalities of i -flats	the dimensions of i -flats
t -funtion	the number of j -flats between an i -flat and a k -flat	???
flats and design	If flats have all of the subsets whose cardinalities are less than $t-1, m$ -flats are t -design ($m \geq t$)	??? (main result)


	PMD	q -PMD
How to construct non trivial (q)-PMD	<ul style="list-style-type: none"> • Projective geometries • Affine geometries • Affine triple systems • Steiner system 	<ul style="list-style-type: none"> • Steiner system • ??? • ??? • ???
α -sequence	the cardinalities of i -flats	the dimensions of i -flats
t -funtion	the number of j -flats between an i -flat and a k -flat	???
α -sequence	If flats have all of the subsets whose cardinalities are less than $t-1, m$ -flats are t -design ($m \geq t$)	??? (main result)

Definition [q -PMD]

A q -perfect matroid design (q -PMD) is a q -matroid with the property that any two flats of the same rank have the same dimension.

Definition [α -sequence]

- α_i : the dimension of the i -flats of a q -PMD
- $\{\alpha_i\}_{i=0}^{r(E)}$: is called an α -**sequence** of the q -PMD



Example 1


$U_{k,n}[\mathbb{F}_q]$ is a q -PMD with an α -sequence $(0, 1, \dots, k-1, n)$.

\therefore All of i -dimensional subspaces are i -flats of $U_{k,n}[\mathbb{F}_q]$ ($i \leq k-1$). The ground space is the k -flat.

Example 2

If the ground set \mathbb{F}_q^{lm} is partitioned into \mathbb{F}_q^m , the partition induces a q -PMD with an α -sequence $(0, m, lm)$.

2-flat



	PMD	q -PMD
How to construct non trivial (q)-PMD	<ul style="list-style-type: none"> • Projective geometries • Affine geometries • Affine triple systems • Steiner system 	<ul style="list-style-type: none"> • Steiner system • ??? • ??? • ???
α -sequence	the cardinalities of i -flats	the dimensions of i -flats
t -funtion	the number of j -flats between an i -flat and a k -flat	???
flats and design	If flats have all of the subsets whose cardinalities are less than $t-1, m$ -flats are t -design ($m \geq t$)	??? (main result)

	PMD	q -PMD
How to construct non trivial (q)-PMD	<ul style="list-style-type: none"> • Projective geometries • Affine geometries • Affine triple systems • Steiner system 	<ul style="list-style-type: none"> • Steiner system • ??? • ??? • ???
α -sequence	the cardinalities of i -flats	the dimensions of i -flats
t -funtion	the number of j -flats between an i -flat and a k -flat	???
flats and design	If flats have all of the subsets whose cardinalities are less than $t-1, m$ -flats are t -design ($m \geq t$)	??? (main result)

- $F_i \in \mathcal{F}_i, F_k \in \mathcal{F}_k$ with $F_i \leq F_k$

Proposition

The number $|\mathcal{F}_j(F_i, F_k)|$ of j -flat F_j with $F_i \leq F_j \leq F_k$ is independent of the choice of F_i and F_k .


Definition [t -function of q-PMD]

We define t -function of M as follows:

$$t_M(i, j, k) := |\mathcal{F}_j(F_i, F_k)|.$$


Proposition

The number $|\mathcal{F}_j(F_i, F_k)|$ of j -flat F_j with $F_i \leq F_j \leq F_k$ is independent of the choice of F_i and F_k .



Proposition

The number $|\mathcal{F}_j(F_i, F_k)|$ of j -flat F_j with $F_i \leq F_j \leq F_k$ is independent of the choice of F_i and F_k .



t -function is calculated by α -sequence

Proposition

Let $M = (E, r)$ be a q -PMD with α -sequence $\{\alpha_i\}_{i=0}^{i=r(E)}$ and t -function t_M . Then,

$$t_M(i, j, k) = \prod_{l=0}^{j-i-1} \frac{q^{\alpha_k} - q^{\alpha_{i+l}}}{q^{\alpha_j} - q^{\alpha_{i+l}}}$$

holds.

	PMD	q -PMD
How to construct non trivial (q)-PMD	<ul style="list-style-type: none"> • Projective geometries • Affine geometries • Affine triple systems • Steiner system 	<ul style="list-style-type: none"> • Steiner system • ??? • ??? • ???
α -sequence	the cardinalities of i -flats	the dimensions of i -flats
t -funtion	the number of j -flats between an i -flat and a k -flat	the function returns the number of j -flats between an i -flat and a k -flat
flats and design	If flats have all of the subsets whose cardinalities are less than $t-1, m$ -flats are t -design ($m \geq t$)	??? (main result)

	PMD	q -PMD
How to construct non trivial (q)-PMD	<ul style="list-style-type: none"> • Projective geometries • Affine geometries • Affine triple systems • Steiner system 	<ul style="list-style-type: none"> • Steiner system • ??? • ??? • ???
α -sequence	the cardinalities of i -flats	the dimensions of i -flats
t -funtion	the number of j -flats between an i -flat and a k -flat	the function returns the number of j -flats between an i -flat and a k -flat
flats and design	If flats have all of the subsets whose cardinalities are less than $t-1, m$ -flats are t -design ($m \geq t$)	??? (main result)


i -Flats of the q -PMD are the blocks of a t -design.

- integer $t : 0 \leq t \leq n$


Theorem

If $\begin{bmatrix} E \\ i \end{bmatrix}_q \subset \mathcal{F}$ for all integers i satisfying $0 \leq i \leq t - 1$, then (E, \mathcal{F}_k) is a subspace design $t\text{-}(n, \alpha_k, t_M(t, k, r(E)))$ for all integers k satisfying $t \leq k \leq r(E)$


sketch of proof




sketch of proof




sketch of proof




sketch of proof



sketch of proof




sketch of proof




(qF3) $\forall F \in \mathcal{F}, \forall x \in \begin{bmatrix} E \\ 1 \end{bmatrix}_q \setminus \begin{bmatrix} F \\ 1 \end{bmatrix}_q, \exists! F' \in \mathcal{F}$ with $r(F') = r(F) + 1$
 $s.t. F + x \subseteq F'$.


————— inclusion relation




sketch of proof




sketch of proof






(qF3) $\forall F \in \mathcal{F}, \forall x \in \begin{bmatrix} E \\ 1 \end{bmatrix}_q \setminus \begin{bmatrix} F \\ 1 \end{bmatrix}_q, \exists! F' \in \mathcal{F}$ with $r(F') = r(F) + 1$
s.t. $F + x \subseteq F'$.



sketch of proof



Theorem

If $\begin{bmatrix} E \\ i \end{bmatrix}_q \subset \mathcal{F}$ for all integers i satisfying $0 \leq i \leq t - 1$, then (E, \mathcal{F}_k) is a subspace design $t\text{-}(n, \alpha_k, t_M(t, k, r(E)))$ for all integers k satisfying $t \leq k \leq r(E)$
if $k = 1, t_M(t, t, r(E)) = 1$

Corollary

If $\begin{bmatrix} E \\ i \end{bmatrix}_q \subset \mathcal{F}$ for all integers i satisfying $0 \leq i \leq t - 1$, then (E, \mathcal{F}_t) is a q -Steiner system $\mathcal{S}(t, \alpha_t, n; q)$

Theorem

M is induced by q -Stienr system $\mathcal{S}(t, k, n)$

E. Byrn and others 2022+




Corollary

an α -sequence of M is $(0, 1, \dots, t - 1, k, n)$

Theorem

M is induced by q -Steiner system $\mathcal{S}(t, k, n)$

E. Byrn and others 2022+

 **Corollary** an α -sequence of M is $(0, 1, \dots, t - 1, k, n)$

sketch of proof

$$\begin{aligned} |\mathcal{F}_j| &= t_M(0, j, r(E)) = \prod_{l=0}^{j-1} \frac{q^n - q^l}{q^j - q^l} = \begin{bmatrix} n \\ j \end{bmatrix}_q \quad \left(\because t_M(i, j, k) = \prod_{l=0}^{j-i-1} \frac{q^{\alpha_k} - q^{\alpha_{i+l}}}{q^{\alpha_j} - q^{\alpha_{i+l}}} \right) \\ \therefore \mathcal{F}_j &= \begin{bmatrix} E \\ j \end{bmatrix}_q \quad (0 \leq j \leq t-1) \end{aligned}$$

$\Rightarrow (E, \mathcal{F}_t)$ is a q -Steiner system $\mathcal{S}(t, k, n; q)$ (∴ Corollary)

	PMD	q -PMD
How to construct non trivial (q)-PMD	<ul style="list-style-type: none"> • Projective geometries • Affine geometries • Affine triple systems • Steiner system 	<ul style="list-style-type: none"> • Steiner system • ??? • ??? • ???
α -sequence	the cardinalities of i -flats	the dimensions of i -flats
t -funtion	the number of j -flats between an i -flat and a k -flat	the function returns the number of j -flats between an i -flat and a k -flat
flats and design	If flats have all of the subsets whose cardinalities are less than $t-1, m$ -flats are t -design ($m \geq t$)	If flats have all of the subsets whose dimensions are less than $t-1$, m -flats are t -subspace design ($m \geq t$)

Problems for q -PMD

- Are there any other non-trivial q -Steiner systems?
 - A q -Steiner system induce a q -PMD[1]
 - The only known q -Steiner system is $\mathcal{S}(2, 3, 13; 2)$ [3]
- Are there q -PMDs not induced by q -Steiner systems?
 - There are some PMDs not induced by Steiner system
 - Projective geometries
 - Affine geometries
 - Affine triple systems

[1] E. Byrne et al. : Constructions of new Mtroids and Designs over \mathbb{F}_q , (2022)

[2] M. Deza

[3] M. Braun et al.: EXISTENCE OF q -ANALOGS OF STEINER SYSTEMS (2013)

	PMD	q -PMD
How to construct non trivial (q)-PMD	<ul style="list-style-type: none"> • Projective geometries • Affine geometries • Affine triple systems • Steiner system 	<ul style="list-style-type: none"> • Steiner system • ??? • ??? • ???
α -sequence	the cardinalities of i -flats	the dimensions of i -flats
t -funtion	the number of j -flats between an i -flat and a k -flat	the function returns the number of j -flats between an i -flat and a k -flat
flats and design	If flats have all of the subsets whose cardinalities are less than $t-1$, m -flats are t -design ($m \geq t$)	If flats have all of the subsets whose dimensions are less than $t-1$, m -flats are t -subspace design ($m \geq t$)

Matroid is a pair (E, r) satisfies (R1), (R2) and (R3).

Definition 1.1. [matroid]

A q -matroid M is a pair (E, r) where r is an integer-valued function defined on $\mathcal{L}(E)$ with the following properties:

$$(R1) \quad \forall A \subseteq E, 0 \leq r(A) \leq |A|.$$

$$(R2) \quad \forall A, B \subseteq E \text{ with } A \subseteq B, 0 \leq r(A) \leq |A|.$$

$$(R3) \quad \forall A, B \subseteq E, r(A \cup B) + r(A \cap B) \leq r(A) + r(B)$$

Remark 1.2. the function r is called **rank function**.

- M : q -PMD

M is induced by q -Stienr system $\mathcal{S}(t, k, n)$

E. Byrn and others 2022+



Theorem

α -sequence of M is $(0, 1, \dots t - 1, k, n)$

sketch of proof

$$\begin{aligned} |\mathcal{F}_j| = t_M(0, j, r(E)) &= \prod_{l=0}^{j-1} \frac{q^n - q^l}{q^j - q^l} = \begin{bmatrix} n \\ j \end{bmatrix}_q \quad \left(\because t_M(i, j, k) = \prod_{l=0}^{j-i-1} \frac{q^{\alpha_k} - q^{\alpha_{i+l}}}{q^{\alpha_j} - q^{\alpha_{i+l}}} \right) \\ \therefore \mathcal{F}_j &= \begin{bmatrix} E \\ j \end{bmatrix}_q \quad (0 \leq j \leq t-1) \end{aligned}$$


Thank you for your kind attention.

- **PMD**

- **α -sequences**
 - represents the numbers of elements in one i-flat
- **t -functions**
 - determine the number of i-flats of q-PMD
- **flats and design**
 - in some conditions, flats are the block of t-design

- **q -PMD**

- **α -sequences**
 - represents the dimension of one i-flat
- **t -functions**
 - determine the number of i-flats of q-PMD
- **flats and t -subspace design**
 - in some conditions, flats are the blocks of a t -subspace design



q -Matroid is a pair (E, r) satisfies (qR1), (qR2) and (qR3).

Definition 1.1. [q -matroid]

$$(qR1) \quad \forall A \in \mathcal{L}(E), \quad 0 \leq r(A) \leq \dim A.$$

$$(qR2) \quad \forall A, B \in \mathcal{L}(E) \text{ with } A \subseteq B, \quad r(A) \leq r(B).$$

$$(qR3) \quad \forall A, B \in \mathcal{L}(E), \quad r(A + B) + r(A \cap B) \leq r(A) + r(B)$$

Remark 1.5. The function r is called **rank function** of M .

Remark 1.5.

We denote \mathcal{F} as the set of flats.

- If a flat F satisfies $r(F) = i$, F is called **i -flat**.
- The collection of i -flat is denoted by \mathcal{F}_i .

Theorem (E. Byrn et al. 2022+)

- (E, \mathcal{B}) : q -Steiner system $\mathcal{S}(t, k, n)$

Let \mathcal{F} be defined as follows:

$$\mathcal{F} := \left\{ \bigcap_{B \in S} B \mid S \subseteq \mathcal{B} \right\}.$$


Then, there exists a q -PMD whose ground set is E and the flats are \mathcal{F} .

Remark

- The α -sequence is $(0, 1, \dots, t-1, k, n)$.

	q -matroid	matroid
1 st axiom	(qF1) $E \in \mathcal{F}$	(qF1) $E \in \mathcal{F}$
2 nd axiom	(qF2) $F_1, F_2 \in \mathcal{F} \Rightarrow F_1 \cap F_2 \in \mathcal{F}$	(qF2) $F_1, F_2 \in \mathcal{F} \Rightarrow F_1 \cap F_2 \in \mathcal{F}$
3 rd axiom	(qF3) $F \in \mathcal{F}, x \in \begin{bmatrix} E \\ 1 \end{bmatrix}_q \setminus \begin{bmatrix} F \\ 1 \end{bmatrix}_q$ $\Rightarrow \exists! F' \in \mathcal{F}_{r(F)+1} \quad s.t. \quad F + x \subseteq F'$	(qF3) $F \in \mathcal{F}, x \in E - X$ $\Rightarrow \exists! F' \in \mathcal{F}_{r(F)+1} \quad s.t. \quad F \cup x \subseteq F'$

Example3



proof

By (qF3), for all 1 dimennal subspace x of F_k , there are unique $i + 1$ flat that includs x and F_i . Now, we count the number of the pair of $i + 1$ -flat and 1 dimensional subspace of them by 2 ways.

We can prove these identities by following the way P. Young and J. Edmond did for normal PMD.

Lemma [Properties of t -function]

Let $M = (E, r)$ be a q -PMD with the t -function t_M . The followings hold:

$$(T0) \quad t_M(i, i, k) = 1, \text{ for } 0 \leq i \leq k \leq r(E).$$

$$(T1) \quad t_M(0, 1, i+1) > t_M(0, 1, i), \text{ for } 0 \leq i \leq r(E) - 1.$$

$$(T2) \quad t_M(i, i+1, k) = \frac{t_M(0, 1, k) - t_M(0, 1, i)}{t_M(0, 1, i+1) - t_M(0, 1, i)}, \text{ for } 0 \leq i < k \leq r(E).$$

$$(T3) \quad t_M(i, j, k) = \frac{t_M(i, l, k) t_M(l, j, k)}{t_M(i, l, j)}, \text{ for } 0 \leq i \leq l \leq j \leq k \leq r(E).$$

proof

By (qF3), for all 1 dimennal subspace x of F_k , there are unique $i + 1$ flat that includs x and F_i . Now, we count the number of the pair of $i + 1$ -flat and 1 dimensional subspace of them. We define

$$C := \{(F_{i+1}, a) \mid F_{i+1} \in \mathcal{F}_{i+1}(F_i, F_k) \wedge a \in \begin{bmatrix} F_{i+1} \\ 1 \end{bmatrix}_q\}.$$

Let $t = |\mathcal{F}_{i+1}(F_i, F_k)|$. then $|C| = t \begin{bmatrix} \alpha_k \\ 1 \end{bmatrix}_q$


$r(X) = \min(|X|, k)$ is an example of rank function.


Example1.3. [The uniform matroid]


Let E be a finite set with n elements and k be a integer satisfying $0 \leq k \leq n$. We define the rank function r as follow:


$$r(X) = \min(|X|, k).$$


Then, the pair (E, r) is a matroid.


$(qF1)$ 

$(qF2)$ 

$(qF2)$ 

$(qF3)$ 

$(qF3)$ 

$(qF3)$ 

proof

$$\left[\begin{array}{ccccc} 1 & 1 & \cdots & 1 & 1 \\ 1 & 1 & \cdots & 1 & 1 \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 1 & 1 & \cdots & 1 & 1 \\ 1 & 0 & \cdots & 0 & 0 \\ 1 & 0 & \cdots & 0 & 0 \\ 0 & 1 & \cdots & 0 & 0 \\ 0 & 1 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & 0 & 1 \\ 0 & 0 & \cdots & 0 & 1 \end{array} \right] \quad \left. \begin{array}{l} \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \end{array} \right\} \text{1-dimensional subspaces of } F_k. \\ \left. \begin{array}{c} \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \end{array} \right\} i+1 \text{ flat in } \mathcal{F}_{i+1}(F_i, F_k).$$

proof

$$F_{i+1}$$

x	$\begin{bmatrix} 1 & 1 & \cdots & 1 & 1 \\ 1 & 1 & \cdots & 1 & 1 \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 1 & 1 & \cdots & 1 & 1 \\ 1 & 0 & \cdots & 0 & 0 \\ 1 & 0 & \cdots & 0 & 0 \\ 0 & 1 & \cdots & 0 & 0 \\ 0 & 1 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & 0 & 1 \\ 0 & 0 & \cdots & 0 & 1 \end{bmatrix}$	
		<p>1-dimensional subspaces of F_k.</p>

$i+1$ flat in $\mathcal{F}_{i+1}(F_i, F_k)$.

$$1 \cdots x \subseteq F_{i+1}$$

$$0 \cdots x \not\subseteq F_{i+1}$$

proof

$$F_{i+1}$$

x	$\begin{bmatrix} 1 & 1 & \cdots & 1 & 1 \\ 1 & 1 & \cdots & 1 & 1 \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 1 & 1 & \cdots & 1 & 1 \\ 1 & 0 & \cdots & 0 & 0 \\ 1 & 0 & \cdots & 0 & 0 \\ 0 & 1 & \cdots & 0 & 0 \\ 0 & 1 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & 0 & 1 \\ 0 & 0 & \cdots & 0 & 1 \end{bmatrix}$	
		

$i+1$ flat in $\mathcal{F}_{i+1}(F_i, F_k)$.

$$1 \cdots x \subseteq F_{i+1}$$

$$0 \cdots x \not\subseteq F_{i+1}$$

proof

1	1	...	1	1
1	1	...	1	1
:	:	:	:	:
1	1	...	1	1
1	0	...	0	0
1	0	...	0	0
0	1	...	0	0
0	1	...	0	0
:	:	:	:	:
0	0	...	0	1
0	0	...	0	1

The number of 1

$i+1$ flat in $\mathcal{F}_{i+1}(F_i, F_k)$

proof

$$\begin{bmatrix} 1 & 1 & \cdots & 1 & 1 \\ 1 & 1 & \cdots & 1 & 1 \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 1 & 1 & \cdots & 1 & 1 \\ 1 & 0 & \cdots & 0 & 0 \\ 1 & 0 & \cdots & 0 & 0 \\ 0 & 1 & \cdots & 0 & 0 \\ 0 & 1 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & 0 & 1 \\ 0 & 0 & \cdots & 0 & 1 \end{bmatrix}$$


$i+1$ flat in $\mathcal{F}_{i+1}(F_i, F_k)$

The number of 1

$$\left[\begin{array}{c} \alpha_{i+1} \\ 1 \end{array} \right]_a$$

proof

1	1	...	1	1
1	1	...	1	1
:	:	:	:	:
1	1	...	1	1
1	0	...	0	0
1	0	...	0	0
0	1	...	0	0
0	1	...	0	0
:	:	:	:	:
0	0	...	0	1
0	0	...	0	1

i+1 flat in $\mathcal{F}_{i+1}(F_i, F_k)$

The number of 1

$$|\mathcal{F}_{i+1}(F_i, F_k)| \begin{bmatrix} \alpha_{i+1} \\ 1 \end{bmatrix}_q$$

proof

1	1	...	1	1
1	1	...	1	1
:	:	:	:	:
1	1	...	1	1
1	0	...	0	0
1	0	...	0	0
0	1	...	0	0
0	1	...	0	0
:	:	:	:	:
0	0	...	0	1
0	0	...	0	1

i+1 flat in $\mathcal{F}_{i+1}(F_i, F_k)$

1 dimensional
subspace of F_k

proof

1	1	...	1	1
1	1	...	1	1
:	:	:	:	:
1	1	...	1	1
1	0	...	0	0
1	0	...	0	0
0	1	...	0	0
0	1	...	0	0
:	:	:	:	:
0	0	...	0	1
0	0	...	0	1

1 dimensional
subspace of F_k

The number of 1 $|\mathcal{F}_{i+1}(F_i, F_k)| \times \begin{bmatrix} \alpha_i \\ 1 \end{bmatrix}_q +$

proof

$$\begin{bmatrix} 1 & 1 & \cdots & 1 & 1 \\ 1 & 1 & \cdots & 1 & 1 \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 1 & 1 & \cdots & 1 & 1 \\ \hline 1 & 0 & \cdots & 0 & 0 \\ 1 & 0 & \cdots & 0 & 0 \\ 0 & 1 & \cdots & 0 & 0 \\ 0 & 1 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & 0 & 1 \\ 0 & 0 & \cdots & 0 & 1 \end{bmatrix}$$

1 dimensional
subspace of F_k

The number of 1 $|\mathcal{F}_{i+1}(F_i, F_k)| \times \left[\begin{matrix} \alpha_i \\ 1 \end{matrix} \right]_q + \left[\begin{matrix} \alpha_k \\ 1 \end{matrix} \right]_q - \left[\begin{matrix} \alpha_i \\ 1 \end{matrix} \right]_q$

Corollary [t -function]

Let M be a q -PMD with flats \mathcal{F} and F_i, F_j are an i -flat and a k -flat respectively with $F_i \subseteq F_j$.

We define **t -function** of M as follows:

$$t_M(i, j, k) := |\mathcal{F}_j(F_i, F_k)|$$

Flat axiom

Definition 1.7.

Let E be a finite dimension vectorspace over \mathbb{F}_q and $\mathcal{F} \subseteq \mathcal{L}(E)$. We define flat axioms as follows:

$$(qF1) \quad E \in \mathcal{F}.$$

$$(qF2) \quad \forall F_1, F_2 \in \mathcal{F}, \quad F_1 \cap F_2 \in \mathcal{F}.$$

$$(qF3) \quad \forall F \in \mathcal{F}, \quad \forall x \in \begin{bmatrix} E \\ 1 \end{bmatrix}_q \setminus \begin{bmatrix} F \\ 1 \end{bmatrix}_q, \quad \exists! F' \in \mathcal{F} \text{ with } r(F') = r(F) + 1 \\ \text{s.t. } F + x \subseteq F'.$$

proof

$$\left[\begin{array}{ccccc} 1 & 1 & \cdots & 1 & 1 \\ 1 & 1 & \cdots & 1 & 1 \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 1 & 1 & \cdots & 1 & 1 \\ 1 & 0 & \cdots & 0 & 0 \\ 1 & 0 & \cdots & 0 & 0 \\ 0 & 1 & \cdots & 0 & 0 \\ 0 & 1 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & 0 & 1 \\ 0 & 0 & \cdots & 0 & 1 \end{array} \right] \quad \begin{matrix} F_i \\ x \end{matrix}$$

$i+1$ flat in $\mathcal{F}_{i+1}(F_i, F_k)$

The number of 1 $|\mathcal{F}_{i+1}(F_i, F_k)| \times \left[\begin{matrix} \alpha_i \\ 1 \end{matrix} \right]_q + \left[\begin{matrix} \alpha_k \\ 1 \end{matrix} \right]_q - \left[\begin{matrix} \alpha_i \\ 1 \end{matrix} \right]_q$

proof

$$\left[\begin{array}{cccc|c} 1 & 1 & \cdots & 1 & 1 \\ 1 & 1 & \cdots & 1 & 1 \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 1 & 1 & \cdots & 1 & 1 \\ 1 & 0 & \cdots & 0 & 0 \\ 1 & 0 & \cdots & 0 & 0 \\ 0 & 1 & \cdots & 0 & 0 \\ 0 & 1 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & 0 & 1 \\ 0 & 0 & \cdots & 0 & 1 \end{array} \right] \quad \begin{matrix} F_i \\ x \end{matrix}$$

$i+1$ flat in $\mathcal{F}_{i+1}(F_i, F_k)$

The number of 1 $|\mathcal{F}_{i+1}(F_i, F_k)| \times \left[\begin{matrix} \alpha_i \\ 1 \end{matrix} \right]_q + \left[\begin{matrix} \alpha_k \\ 1 \end{matrix} \right]_q - \left[\begin{matrix} \alpha_i \\ 1 \end{matrix} \right]_q$

proof

$$\left[\begin{array}{cccc|c} 1 & 1 & \cdots & 1 & 1 \\ 1 & 1 & \cdots & 1 & 1 \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 1 & 1 & \cdots & 1 & 1 \\ 1 & 0 & \cdots & 0 & 0 \\ 1 & 0 & \cdots & 0 & 0 \\ 0 & 1 & \cdots & 0 & 0 \\ 0 & 1 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & 0 & 1 \\ 0 & 0 & \cdots & 0 & 1 \end{array} \right] \quad F_i$$

F_i

x

$i+1$ flat in $\mathcal{F}_{i+1}(F_i, F_k)$

$$\text{The number of 1} \dots \dots |F_{i+1}(F_i, F_k)| \times \left[\begin{matrix} \alpha_i \\ 1 \end{matrix} \right]_q + \left[\begin{matrix} \alpha_k \\ 1 \end{matrix} \right]_q - \left[\begin{matrix} \alpha_i \\ 1 \end{matrix} \right]_q$$

proof

$$\begin{bmatrix} 1 & 1 & \cdots & 1 & 1 \\ 1 & 1 & \cdots & 1 & 1 \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 1 & 1 & \cdots & 1 & 1 \\ \hline 1 & 0 & \cdots & 0 & 0 \\ 1 & 0 & \cdots & 0 & 0 \\ 0 & 1 & \cdots & 0 & 0 \\ 0 & 1 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & 0 & 1 \\ 0 & 0 & \cdots & 0 & 1 \end{bmatrix}$$



1 dimensional
subspace of

The number of 1

$$|\mathcal{F}_{i+1}(F_i, F_k)| \times \left[\begin{matrix} \alpha_i \\ 1 \end{matrix} \right]_q + \left[\begin{matrix} \alpha_k \\ 1 \end{matrix} \right]_q - \left[\begin{matrix} \alpha_i \\ 1 \end{matrix} \right]_q$$

proof

$$\begin{bmatrix} 1 & 1 & \cdots & 1 & 1 \\ 1 & 1 & \cdots & 1 & 1 \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 1 & 1 & \cdots & 1 & 1 \\ 1 & 0 & \cdots & 0 & 0 \\ 1 & 0 & \cdots & 0 & 0 \\ 0 & 1 & \cdots & 0 & 0 \\ 0 & 1 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & 0 & 1 \\ 0 & 0 & \cdots & 0 & 1 \end{bmatrix}$$

$$|\mathcal{F}_{i+1}(F_i, F_k)| \times \left[\begin{smallmatrix} \alpha_{i+1} \\ 1 \end{smallmatrix} \right]_q = |\mathcal{F}_{i+1}(F_i, F_k)| \times \left[\begin{smallmatrix} \alpha_i \\ 1 \end{smallmatrix} \right]_q + \left[\begin{smallmatrix} \alpha_k \\ 1 \end{smallmatrix} \right]_q - \left[\begin{smallmatrix} \alpha_i \\ 1 \end{smallmatrix} \right]_q$$

$$\therefore |\mathcal{F}_{i+1}(F_i, F_k)| = \frac{q^{\alpha_k} - q^{\alpha_i}}{q^{\alpha_{i+1}} - q^{\alpha_i}}$$

We can prove these identities by following the way P. Young and J. Edmond did for normal PMD.

Lemma [Properties of t -function]

Let $M = (E, r)$ be a q -PMD with the t -function t_M . The followings hold:

$$(T0) \quad t_M(i, i, k) = 1, \text{ for } 0 \leq i \leq k \leq r(E).$$

$$(T1) \quad t_M(0, 1, i+1) > t_M(0, 1, i), \text{ for } 0 \leq i \leq r(E) - 1.$$

$$(T2) \quad t_M(i, i+1, k) = \frac{t_M(0, 1, k) - t_M(0, 1, i)}{t_M(0, 1, i+1) - t_M(0, 1, i)}, \text{ for } 0 \leq i < k \leq r(E).$$

$$(T3) \quad t_M(i, j, k) = \frac{t_M(i, l, k) t_M(l, j, k)}{t_M(i, l, j)}, \text{ for } 0 \leq i \leq l \leq j \leq k \leq r(E).$$

$$(T3) \quad t_M(i, j, k) = \frac{t_M(i, l, k)t_M(l, j, k)}{t_M(i, l, j)}, \text{ for } 0 \leq i \leq l \leq j \leq k \leq r(E).$$

$$t_M(i, j, k) = \frac{t_M(i, i+1, k)}{t_M(i, i+1, j)} t_M(i+1, j, k)$$

$$(T3) \quad t_M(i, j, k) = \frac{t_M(i, l, k)t_M(l, j, k)}{t_M(i, l, j)}, \text{ for } 0 \leq i \leq l \leq j \leq k \leq r(E).$$

$$\begin{aligned} t_M(i, j, k) &= \frac{t_M(i, i+1, k)}{t_M(i, i+1, j)} t_M(i+1, j, k) \\ &= \frac{t_M(i, i+1, k)}{t_M(i, i+1, j)} \cdot \frac{t_M(i+1, i+2, k)}{t_M(i+1, i+2, j)} \cdot t_M(i+2, j, k) \\ &\vdots \\ &= \prod_{l=0}^{j-i-1} \frac{t_M(i+l, i+l+1, k)}{t_M(i+l, i+l+1, j)} \end{aligned}$$

by Lemma


$$t_M(i, i+1, k) := |\mathcal{F}_{i+1}(F_i, F_k)| = \frac{q^{\alpha_k} - q^{\alpha_i}}{q^{\alpha_{i+1}} - q^{\alpha_i}}$$


$$\begin{aligned} t_M(i, j, k) &= \frac{t_M(i, i+1, k)}{t_M(i, i+1, j)} t_M(i+1, j, k) \\ &= \frac{t_M(i, i+1, k)}{t_M(i, i+1, j)} \cdot \frac{t_M(i+1, i+2, k)}{t_M(i+1, i+2, j)} \cdot t_M(i+2, j, k) \\ &\vdots \\ &= \prod_{l=0}^{j-i-1} \frac{t_M(i+l, i+l+1, k)}{t_M(i+l, i+l+1, j)} \\ &= \prod_{l=0}^{j-i-1} \frac{q^{\alpha_k} - q^{\alpha_{i+l}}}{q^{\alpha_{i+j+1}} - q^{\alpha_{i+l}}} \cdot \frac{q^{\alpha_{i+j+1}} - q^{\alpha_{i+l}}}{q^{\alpha_j} - q^{\alpha_{i+l}}} = \prod_{l=0}^{j-i-1} \frac{q^{\alpha_k} - q^{\alpha_{i+l}}}{q^{\alpha_j} - q^{\alpha_{i+l}}} \end{aligned}$$

Known q -PMDs.

α -sequence	$\mathcal{F}_{r(E)-1}$	description
$(0, 1, \dots, k-1, n)$	$\mathcal{S}(k-1, k-1, n; q)$	uniform q -matroid $U_{k,n}[\mathbb{F}_q]$
$(0, m, lm)$	$\mathcal{S}(1, m, lm; q)$	split of vector space
$(0, 1, 3, 13)$	$\mathcal{S}(2, 3, 13; 2)$	only known non-trivial q -PMD [1] \mathbb{F}_q

寸法や色は場合に応じて調整しましょう





3つあることが一目でわかる

カード①

カードの見出し


カード②

カードの見出し

カード③

カードの見出し

順序関係がある場合は並列よりこちらが良い



カードの間隔やアスペクト比は適宜調整

カード①

カードの見出し

カード③

カードの見出し

カード②

カードの見出し

カード④

カードの見出し

循環がある場合はこんな感じが良い

